

Code	Level of MDM (Based on 2 out of 3 Elements of MDM)	Elements of Medical Decision Making		
		Number and Complexity of Problems Addressed	Amount and/or Complexity of Data to be Reviewed and Analyzed <i>*Each unique test, order, or document contributes to the combination of 2 or combination of 3 in Category 1 below.</i>	Risk of Complications and/or Morbidity or Mortality of Patient Management
99211	N/A	N/A	N/A	N/A
99202 99212	Straightforward	Minimal • 1 self-limited or minor problem	Minimal or none	Minimal risk of morbidity from additional diagnostic testing or treatment
99203 99213	Low	Low • 2 or more self-limited or minor problems; or • 1 stable chronic illness; or • 1 acute, uncomplicated illness or injury	Limited <i>(Must meet the requirements of at least 1 of the 2 categories)</i> Category 1: Tests and documents • Any combination of 2 from the following: • Review of prior external note(s) from each unique source*; • review of the result(s) of each unique test*; • ordering of each unique test* or Category 2: Assessment requiring an independent historian(s) <i>(For the categories of independent interpretation of tests and discussion of management or test interpretation, see moderate or high)</i>	Low risk of morbidity from additional diagnostic testing or treatment
99204 99214	Moderate	Moderate • 1 or more chronic illnesses with exacerbation, progression, or side effects of treatment; or • 2 or more stable chronic illnesses; or • 1 undiagnosed new problem with uncertain prognosis; or • 1 acute illness with systemic symptoms; or • 1 acute complicated injury	Moderate <i>(Must meet the requirements of at least 1 out of 3 categories)</i> Category 1: Tests, documents, or independent historian(s) • Any combination of 3 from the following: • Review of prior external note(s) from each unique source*; • Review of the result(s) of each unique test*; • Ordering of each unique test*; • Assessment requiring an independent historian(s) or Category 2: Independent interpretation of tests • Independent interpretation of a test performed by another physician/other qualified health care professional (not separately reported); or Category 3: Discussion of management or test interpretation • Discussion of management or test interpretation with external physician/other qualified health care professional/appropriate source (not separately reported)	Moderate risk of morbidity from additional diagnostic testing or treatment <i>Examples only:</i> • Prescription drug management • Decision regarding minor surgery with identified patient or procedure risk factors • Decision regarding elective major surgery without identified patient or procedure risk factors • Diagnosis or treatment significantly limited by social determinants of health
99205 99215	High	High • 1 or more chronic illnesses with severe exacerbation, progression, or side effects of treatment; or • 1 acute or chronic illness or injury that poses a threat to life or bodily function	Extensive <i>(Must meet the requirements of at least 2 out of 3 categories)</i> Category 1: Tests, documents, or independent historian(s) • Any combination of 3 from the following: • Review of prior external note(s) from each unique source*; • Review of the result(s) of each unique test*; • Ordering of each unique test*; • Assessment requiring an independent historian(s) or Category 2: Independent interpretation of tests • Independent interpretation of a test performed by another physician/other qualified health care professional (not separately reported); or Category 3: Discussion of management or test interpretation • Discussion of management or test interpretation with external physician/other qualified health care professional/appropriate source (not separately reported)	High risk of morbidity from additional diagnostic testing or treatment <i>Examples only:</i> • Drug therapy requiring intensive monitoring for toxicity • Decision regarding elective major surgery with identified patient or procedure risk factors • Decision regarding emergency major surgery • Decision regarding hospitalization • Decision not to resuscitate or to de-escalate care because of poor prognosis

Source: The American Medical Association